# Advancing Data Assimilation as a Science Hub Weather prediction and beyond



Takemasa Miyoshi

Ph.D. (Meteorology)
Data Assimilation Scientist

Data Assimilation Research Team

**RIKEN** 



#### Who am I?

B.S. from Kyoto U JMA administration (2y) JMA NWP (1.25y) UMD (2y, M.S. and Ph.D.) **JMA NWP (3.5y)** UMD (4y) RIKEN (5.5y+)

#### http://data-assimilation.riken.jp/~miyoshi/

#### Takemasa Miyoshi, Ph.D.

Team Leader

<u>Data Assimilation Research Team</u> RIKEN Center for Computational Science

#### Deputy Director

RIKEN interdisciplinary Theoretical and Mathematical Sciences
(iTHEMS) Program

Chief Scientist
Prediction Science Laboratory
RIKEN Cluster for Pioneering Research

#### Visiting Professor

University of Maryland, College Park

#### Affiliate Professor

Graduate School of Science, Kyoto University

#### Visiting Principal Scientist

Application Laboratory, JAMSTEC

#### Research Counselor

<u>Servicio Meteorológico Nacional (National Meteorological Service),</u>
<u>Argentina</u>



#### **Education**

- 2005 Ph.D. in Meteorology, University of Maryland, College Park, Maryland, USA (Dissertation PDF)
- 2004 M.S. in Meteorology, University of Maryland, College Park, Maryland, USA (Scholarly Paper PDF)
- 2000 B.S. in Physics, Faculty of Science, Kyoto University, Kyoto, Japan



#### RIKEN Center for Computational Science (R-CCS)

## Japan's flagship institute for computational science

#### **Missions:**

- 1) Development & operation of the Japanese flagship supercomputer
  - 2) Center of Excellence for research on computational science



## Japan's flagship institute for computational science

#### Missions:

- 1) Development & operation of the Japanese flagship supercomputer
  - 2) Center of Excellence for research on computational science






#### **Data Assimilation Research Team**

Data Assimilation Research Team was launched in October, 2012, in RIKEN Advanced Institute for Computational Science (AICS), conveniently located in the beautiful and historic city of Kobe. <u>RIKEN</u> is known as the flagship research institution in Japan. On April 1, 2018, RIKEN AICS was renamed <u>RIKEN Center for Computational Science (R-CCS)</u>. R-CCS is operating the world's leading K computer, and also has a strong Research Division. R-CCS takes the lead in advancing the computational science and aims to be an international center of excellence for computational science in collaboration with a wide range of research organizations. R-CCS integrates the computer science and computational science to conduct most advanced research and development of a wide range of applied scientific computation, as well as of high performance computing technologies.

Data assimilation is a cross-disciplinary science to synergize numerical simulations and observational data, using statistical methods and applied mathematics. As computers become more powerful and enable more precise simulations, it will become more important to compare the simulation with actual observations.

Data Assimilation Research Team ("DA team") performs cutting-edge research and development on advanced data assimilation methodsand their rapplications, aiming at integrating computer simulations and observational data in the wisest way. Particularly, the DA team will tackle challenging problems of developing efficient and accurate data assimilation systems for high-dimensional simulations with large amount of data. The specific are include 1) research on parallel-efficient algorithms for data assimilation with the super-parallel K computer, 2) research on data assimilation metho and applications by taking advantage of the world-leading K computer, and 3) development of most advanced data assimilation software optimized the K computer.



February 21, 2018 at R-CCS



#### <u> http://www.data-assimilation.riken.jp/</u>





Guo-Yuan Lien





Research


Associate



## Data Assimilation Research Team To be a world's leader in DA research







#### Core

DA theory

Algorithm

Computation

#### Lead

Leading research in NWP


Toward the world's top level

#### Pioneer

Cross-cutting research beyond NWP

Broader applications

### Global 870-m simulation (Miyamoto et al. 2013)



©JAMSTEC • AORI (SPIRE Field3), RIKEN/AICS Visualized by Ryuji Yoshida





cf. TEDxSannomiya

©JAMSTEC • AORI (SPIRE Field3), RIKEN/AICS Visualized by Ryuji Yoshida

http://tedxsannomiya.com/speakers/takemasa-miyoshi/



About RIKEN

Research

News & Media

Careers

Outreach

Community



Home > News & Media > Press Releases > Press Releases 2018 >



#### Himawari-8 data assimilated simulation enables 10-minute updates of rain and flood predictions

Using the power of Japan's K computer, scientists from the RIKEN Advanced Institute for Computational Science and collaborators have shown that incorporating satellite data at frequent intervals-ten minutes in the case of this study-into weather prediction models can significantly improve the rainfall predictions of the models and allow more precise predictions of the rapid development of a typhoon.

Weather prediction models attempt to predict future weather by running simulations based on current conditions taken from various sources of data. However, the inherently complex nature of the systems, coupled with the lack of precision and timeliness of the data, makes it difficult to conduct accurate predictions, especially with weather systems such as sudden precipitation.

As a means to improve models, scientists are using powerful supercomputers to run simulations based on more frequently updated and accurate data. The team led by Takemasa Miyoshi of AICS decided to work with data from Himawari-8, a geostationary satellite that began operating in 2015. Its instruments can scan the entire area it covers every ten minutes in both visible and infrared light, at a resolution of up to 500 meters, and the data is provided to meteorological agencies. Infrared measurements are useful for indirectly gauging rainfall, as they make it possible to see where clouds are located and at what altitude.

News & Media	
Press Center	>
Press Releases	~
2018	
2017	>
2016	>
2015	>
2014	>
2013	>
2012	>
2011	>
2010	>
2009	>
2008	>
2007	>
2006	>
2005	>
News	>
Events & Symposiums	>



About RIKEN

Research

News & Media

Careers

↑ Index

→ Next

Outreach

Community



← Previous

Home > News & Media > Press Releases > Press Releases 2018 >

Press Release Like 2 January 18, 2018

#### Himawari-8 data assimilated simulation enables 10-minute updates of rain and flood predictions

Using the power of Japan's K computer, scientists and collaborators have shown that incorporating s this study-into weather prediction models can sig allow more precise predictions of the rapid devel

Weather prediction models attempt to predict future taken from various sources of data. However, the in precision and timeliness of the data, makes it diffic systems such as sudden precipitation.

As a means to improve models, scientists are using powerful supercomputers to run simulations based on more frequently updated and accurate data. The team led by Takemasa Miyoshi of AICS decided to work with data from Himawari-8, a geostationary satellite that began operating in 2015. Its instruments can scan the entire area it covers every ten minutes in both visible and infrared light, at a resolution of up to 500 meters, and the data is provided to meteorological agencies. Infrared measurements are useful for indirectly gauging rainfall, as they make it possible to see where clouds are located and at what altitude.



#### Himawari-8 DA papers published (*Honda et al. 2018a,b*, JGR & MWR)

#### **@AGU**PUBLICATIONS

### JGR

#### Journal of Geophysical Research: Atmospheres

#### **RESEARCH ARTICLE**

10.1002/2017JD027096

#### **Key Points:**

- Himawari-8 satellite observations enable rapid update of precipitation and flood predictions
- Precipitation forecasts are significantly improved by assimilating Himawari-8 all-sky radiances
- Every 10 min refresh with Himawari-8 has the potential to capture the flood risk with longer lead times

Assimilation of Himawari-8 All-Sky Radiances Every 10 Minutes: Impact on Precipitation and Flood Risk Prediction

T. Honda<sup>1</sup> [0], S. Kotsuki<sup>1</sup> [0], G.-Y. Lien<sup>1</sup> [0], Y. Maejima<sup>1</sup>, K. Okamoto<sup>2,1</sup> [0], and T. Miyoshi<sup>1,3,4</sup> [0]

<sup>1</sup>RIKEN Advanced Institute for Computational Science, Kobe, Japan, <sup>2</sup>Meteorological Research Institute, Japan Meteorological Agency, Tsukuba, Japan, <sup>3</sup>Department of Atmospheric and Oceanic Science, University of Maryland, College Park, Maryland, USA, <sup>4</sup>Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan

#### Correspondence to:

T. Honda and T. Miyoshi, takumi.honda@riken.jp; takemasa.miyoshi@riken.jp


### <sup>8</sup>Assimilating All-Sky *Himawari-8* Satellite Infrared Radiances: A Case of Typhoon Soudelor (2015)

TAKUMI HONDA,<sup>a</sup> TAKEMASA MIYOSHI,<sup>b</sup> GUO-YUAN LIEN,<sup>a</sup> SEIYA NISHIZAWA,<sup>a</sup> RYUJI YOSHIDA,<sup>a</sup> SACHIHO A. ADACHI,<sup>a</sup> KOJI TERASAKI,<sup>a</sup> KOZO OKAMOTO,<sup>c</sup> HIROFUMI TOMITA,<sup>a</sup> AND KOTARO BESSHO<sup>d,e</sup>

<sup>a</sup> RIKEN Advanced Institute for Computational Science, Kobe, Japan
<sup>b</sup> RIKEN Advanced Institute for Computational Science, Kobe, Japan, and Department of Atmospheric and Oceanic Science, University of Maryland, College Park, College Park, Maryland, and Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan
<sup>c</sup> Meteorological Research Institute, Japan Meteorological Agency, Tsukuba, and RIKEN Advanced Institute for Computational Science, Kobe, Japan
<sup>d</sup> Meteorological Satellite Center, Japan Meteorological Agency, Kiyose, Japan

(Manuscript received 15 September 2016, in final form 18 November 2017)

#### Himawari-8 DA papers published (*Honda et al. 2018a,b*, JGR & MWR)



#### Correspondence to:

T. Honda and T. Miyoshi, takumi.honda@riken.jp; takemasa.miyoshi@riken.jp

### <sup>8</sup>Assimilating All-Sky *Himawari-8* Satellite Infrared Radiances: A Case of Typhoon Soudelor (2015)

TAKUMI HONDA, TAKEMASA MIYOSHI, GUO-YUAN LIEN, SEIYA NISHIZAWA, RYUJI YOSHIDA, SACHIHO A. ADACHI, KOJI TERASAKI, KOZO OKAMOTO, HIROEUMI TOMITA AND KOTARO BESSHO<sup>d,c</sup>

#### DOI:10.1175/MWR-D-16-0357.1


Atmospheric and Oceanic Science, University of Maryland, College Park, College Park, Maryland,
and Japan Agency for Marine-Earth Science and Technology, Yokohama, Japan

Meteorological Research Institute, Japan Meteorological Agency, Tsukuba, and RIKEN Advanced Institute
for Computational Science, Kobe, Japan

Meteorological Satellite Center, Japan Meteorological Agency, Kiyose, Japan

(Manuscript received 15 September 2016, in final form 18 November 2017)

#### Himawari-8: a new generation geostationary meteorological satellite




16UTC 2 to 13UTC 3 April 2015 MTSAT-2 (VIS) Every 1 hour 16UTC 2 to 13UTC 3 April 2015 Himawari-8 (True Color) Every 10 minutes


(Courtesy of JMA)

### Sources of Big Data



### Sources of Big Data











### **BDA** Big Data Assimilation



### Data Assimilation (DA)



>2

Data-driven ocess-driver tion (DA) Induction Deduction Cyber world Real world Simulations Observations Data Assimilation

>2

### SCALE-LETKF system






Local Ensemble Transform Kalman Filter (*Hunt et al. 2007*)




A Regional Numerical Weather Prediction System developed at RIKEN (Lien et al. 2017, DOI: 10.2151/sola.2017-001)

### Typhoon Soudelor (2015)


• The strongest western north Pacific typhoon in 2015 captured well by Himawari-8




### Himawari-8 impact




### Himawari-8 impact on intensity fcst



### Every 10 min. vs. 30 min. DA




### Intensity forecast (30 min. vs. 10 min.)




Assimilating every 10-min. is essential.

### Another case: September 2015




### Himawari-8 impact



High clouds Brightness Temperature (K) Low clouds/clear sky

### Himawari-8 impact


12-h forecast precipitation (previous 1 h accumulation)



(*Honda et al. 2017*)

12-h forecast rain location much improved


### Pioneering "Big Data Assimilation" Era





### Only in 10 minutes!!

(Courtesy of NICT)





15 EL angles in 5 min)


### Phased Array Weather Radar (PAWR)







(100 m, 100 EL angles in 30 sec)

### Phased Array Radar (every 30 sec.)






### Revolutionary super-rapid 30-sec. cycle



120 times more rapid than hourly update cycles

### 9/11/2014 morning, sudden rain



### 9/11/2014, sudden local rain



© 2016 ZENRIN Image Landsat Image IBCAO Data SIO, NOAA, U.S. Navy, NGA, GEBCO Google earth

# 9/11/2014, sudden local rain

RIKEN Advanced Institute for Computational Science Data Assimilation Research Team

28 14.09. 11 88:0 1:80

# Observation

Simulation (100m Big DA)


>40,000 views #9 of RIKEN channel

Simulation


(TV) (O DA)

Simulation (1km DA)

# 9/11/2014, sudden local rain




## <u>Cyber-Physical fusion</u> for weather prediction Data Assimilation is the key





# Data Assimilation (DA)



Data assimilation best combines real-world data and a model, and brings synergy.









#### Data Assimilation as a tool for vegetation management

Arakida, H., S. Otsuka, S. Kotsuki, T. Miyoshi (RIKEN AICS), S. Moriya (RIKEN CSRS), N. Kobayashi (RIKEN ACCC), J. Aikawa (RIKEN Chief Scientist Laboratories)

# Optimizing an individual-based vegetation model for better prediction of forest evolution.






#### Data assimilation: Application for press forming simulation

Hideyuki Sakamoto, Takemasa Miyoshi (RIKEN AICS)

Masato Takamura (RIKEN Center for Advanced Photonics)





Optimizing press forming conditions by predicting strain to suppress cracks, crumpled, and springback


**Data Assimilation** aims to improve a press forming simulation using measurements of surface strain!



# Industry applications


#### Toward smart hydroelectric dam operations






# Modelling rhythms of the brain individually by a data assimilation approach

Takumi Sase & Keiichi Kitajo (RIKEN BSI)



# Data assimilation in medicine and biology: towards preventive and personalized medicine


Eiryo Kawakami Medical Sciences Innovation Hub Program, RIKEN



#### Forecast Cell Fate by Data Assimilation of Gene Expression Profile

Jun Seita, M.D., Ph.D.

RIKEN MIH / IMS / AIP, Stanford University













#### Development of mathematical model in biology with Approximate Bayesian Computation (ABC)

Tatsuo Shibata (RIKEN Quantitative Biology Center, QBiC)

#### Embryonic patterning is so robust.






Copyright © 2006 Nature Publishing Group Nature Reviews | Molecular Cell Biology

#### Scaled patterning with proportionality

#### Question?

What is the mechanism that underlies the scaling behavior for different size of embryo?




=> What is the mechanism that make the BMP profile independent of embryo size?

### Mathematical model of DV pattern formation

$$\begin{split} \frac{\partial C}{\partial t} &= V_{c} \frac{K_{c}^{h_{c}}}{K_{c}^{h_{c}} + (A+B)^{h_{c}}} - \lambda_{c} \frac{C}{1 + S/K_{c} + (C+BC+AC)/K_{m}} - k_{coolomp}C \cdot B - kC \cdot A + D_{chol}\Delta C \\ \frac{\partial B}{\partial t} &= V_{c} \frac{(A+B)^{h_{c}}}{K_{c}^{h_{c}} + (A+B)^{h_{c}}} - \lambda_{c}B + \lambda_{c} \frac{BC}{1 + S/K_{c} + (C+BC+AC)/K_{m}} - k_{cholomp}C \cdot B + D_{bmp}\Delta B \\ \frac{\partial A}{\partial t} &= V_{c} \frac{K_{c}^{h_{c}}}{K_{c}^{h_{c}} + (A+B)^{h_{c}}} - \lambda_{c}A + \lambda_{c} \frac{AC}{1 + S/K_{c} + (C+BC+AC)/K_{m}} - k_{cholomp}C \cdot A + D_{adinp}\Delta A \\ \frac{\partial S}{\partial t} &= V_{c} \frac{(A+B)^{h_{c}}}{K_{c}^{h_{c}} + (A+B)^{h_{c}}} - \lambda_{c}S + D_{sal}\Delta S \\ \frac{\partial BC}{\partial t} &= -\lambda_{c} \frac{BC}{1 + S/K_{c} + (C+BC+AC)/K_{m}} + k_{cholomp}C \cdot B + D_{cholomp}\Delta BC \\ \frac{\partial AC}{\partial t} &= -\lambda_{c} \frac{AC}{1 + S/K_{c} + (C+BC+AC)/K_{m}} + k_{cholomp}C \cdot A + D_{cholomp}\Delta AC \\ 1311. \end{split}$$

#### Bayesian inference from data




The BMP profile is independent of embryo size

of embryo size Scaling!


#### Crystal growth with data assimilation

Quantum Optodevice Lab., RIKEN M. Jo

# Producing the desired structures based on the growth history data



#### Simulation model of crystal growth



# International Workshop on Uncertainty Quantification Organized by RIKEN Advanced Institute for Computational Science (AICS). Co-organized by RIKEN iTHES/iTHEMS & Graduate School of Science, University of Kyoto. 2018 Feb. 19-21 Venue Auditorium (6F) AICS, RIKEN, Kobe, Japan


#### **Invited Speakers**


- Youssef Marzouk (MIT)
- Omar Knio (KAUST)
- Loic Giraldi (KAUST)
- Olivier Le Maitre (LIMSI-CNRS)
- Pierre Tandeo (IMT-Atlantique)
- Mohamed Iskandarani (University of Miami)
- Serge Guillas (University College London)

7 lecturers + 32 participants from 6 countries









#### http://www.data-assimilation.riken.jp/isda2019/



Scientific Organizing Committee
Chair: Takemasa Miyoshi (RIKEN)
Henry Abarbanel (University of California, San Diego)
Ibrahim Hoteit (KAUST)
Roland Potthast (Deutscher Wetterdienst / University

Roland Potthast (Deutscher Wetterdienst / University of Reading)
Hiromu Seko (JMA,MRI / JAMSTEC)
Martin Weissmann (Ludwig-Maximilians-Universität München)

Serge Guillas (University College London)
Nancy Nichols (University of Reading)
Sebastian Reich (University of Potsdam / University of Reading)
Peter Jan Van Leeuwen (Colorado State University)
Shu-Chih Yang (Taiwan National Central University)

