A Climate Database for the Martian Atmosphere

C. Hourdin, F. Forget, R. Fournier,
F. Hourdin, F. Mathiot and O. Talagrand

Laboratoire de Météorologie Dynamique du CNRS,
Université Paris 6, 4 place Jussieu, 75252 Paris Cedex 05, France

S.R. Lewis, and P.L. Read

University of Oxford, Department of Physics, Clarendon Laboratory,
Parks Road, Oxford OX1 3PU, U.K.

J.-P. Huot

European Space Research and Technology Centre, Noordwijk, The Netherlands

ABSTRACT

A database of statistics which describes the climate of Mars has been constructed by two european teams from Paris (LMD) and Oxford (AOPP), with the support of the European Space Agency (`ESTEC TRP contract 11369/95/NL/JG`). Data are direct outputs of multi-annual simulations performed with the General Circulation Models (GCM) developped jointly at these two laboratories. The models have been validated against the main features of the meteorology of Mars, as observed by spacecraft missions (Mariner 9, Viking, Mars Pathfinder). Primarily designed to provide information about the planetary environment for spacecraft designers (entry profiles in the atmosphere and environment near the surface), this database is also a most usefull tool for the scientific comunity. It is accessible through two interfaces. A fortran software gives access to one dimension profiles with a very sophisticated representation of variability. This version is distributed on a tape. A WWW interface has also been developped:

http://www.lmd.jussieu.fr/mars.html

giving access to means and standard deviations of the variables displayed here after. Data are retrieved either as printable color maps or in various standard formats such as ASCII and NetCDF.

Variable	units	2-D or 3-D
CO_2 ice cover	$kg\,m^{-2}$	2-D
Surface emissivity	none	2-D
Surface temperature	K	2-D
Surface pressure	Pa	2-D
Atmospheric temperature	K	3-D
Zonal (east-west) wind	$m\,s^{-2}$	3-D
Meridional (north-south) wind	$m\,s^{-2}$	3-D
Atmospheric density	$kg\,m^{-3}$	3-D
Boundary layer eddy kinetic energy	$m^2\,s^{-2}$	3-D



Figure 1: *Comparisons between the Martian Climate Database and observations*
Upper: Temperature profiles at the Vinking Lander 1, Viking Lander 2 and Mars Pathfinder sites (solid lines). Each profile is compared with the climatological mean with (dotted lines) and without (dashed lines) the addition of the large-scale and small-scale variability models of the database. The low dust climatology (dotted and dashed lines) is also shown in comparison.
Lower: Observed Mars Pathfinder wind direction as a function of local time of day for the first thirty days of the mission (individual observations are shown as dots, the line is the mean cycle) compared with the MCD mean diurnal cycle for the viking scenario at the same location and time of year (19N, 33W, Ls=142).



Figure 2: *Presentation of the WWW interface*

Here is an example of visualisation of the mean value of temperature in the first layer of the Martian atmosphere. Data are displayed both as a printable color map and as an ASCII file.