

FIRST GLOBAL IMAGES OF MARS FUV DISCRETE AURORA FROM THE EMIRATES MARS MISSION EMUS INSTRUMENT

R. J. Lillis¹, J. Deighan², S. Jain², G. Holsclaw², M. Chaffin², M. Fillingim¹, K. Chirakkil^{2,3}, S. England⁴, D. A. Brain², H. Al Matroushi², F. Lootah⁵, H. Al-Mazmi⁵, S. Xu¹, X. Fang², N. Schneider², J. S. Halekas⁶, S. Ruhunusiri⁶, J. Espley⁷

¹UC Berkeley Space Sciences Laboratory (rlillis@berkeley.edu), ²Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, ³Khalifa University, UAE, ⁴Virginia Institute of Technology, ⁵Mohammed bin Rashid Space Center, Dubai, UAE. ⁶University of Iowa Department of Physics and Astronomy, ⁷NASA Goddard Space Flight Center

Mars Discrete Aurora: current understanding

Three major types of aurora have been discovered to occur in the Mars upper atmosphere: diffuse aurora [1], proton aurora [2], and discrete aurora [3]. The former two are produced by broad energetic charged particle precipitation during extreme solar wind disturbances and direct precipitation of solar wind protons, respectively. The latter is typified by discrete regions of ultraviolet emission from atoms and molecules electronically excited by the precipitation of accelerated electrons into the Martian thermosphere [4, 5]. It has been observed from both disk and limb observations and is most common in regions of strong, topologically open crustal magnetic fields. Aurora observations covering significant fractions of a hemisphere have been observed by MAVEN rarely and only in the mid ultraviolet (> 200 nm) [6].

Data Set: Here we report the first synoptic (or “disk”) observations of Martian discrete aurora in the far ultraviolet (< 200 nm). These are made by the Emirates Mars Ultraviolet Spectrograph (EMUS) on board the Emirates Mars Mission (EMM) and are well-suited to studying aurora due to a) EMM’s high altitude vantage point and regular observation cadence and b) the high sensitivity of the EMUS instrument. The specific auroral observations are from the U-OS2 observation mode designed to study Mars’ inner corona and includes significant coverage of the night side [7], as shown in figure 1.




Figure 1: (left) EMM orbit and typical observation set for U-OS2. Right: box comprised of observation swaths across Mars’ inner corona (< 1.6 Mars radii).

From April 1 until December 31, 2021, 194 swaths provide sufficient coverage of the nightside to study discrete FUV aurora (defined as $> 30\%$ of disk pixels have solar zenith angle $> 110^\circ$).

Spectrum: Emission is strongest at the 130.4 nm oxygen line (3S decay to ground state 3P), though it is also observed in oxygen features at 98.9 nm, 102.7 nm,

and 135.6 nm, as well as possible emission in the CO fourth positive group bands, as shown in the typical spectrum in figure 2.




Figure 2: Typical night side Martian disk spectrum, for both auroral and non-auroral regions.

Results: overlapping swaths can be considered individually or stitched and averaged together to make a single synoptic image, as shown in figure 3. Discrete aurora show patterns broadly matching expectations (i.e. occurring more commonly in strong, open crustal field regions, as in figure 3, left). A notable exception is that the brightest aurora occurred away from crustal fields during the coronal mass ejection impact of July 21, 2021, during which discrete aurora are seen in weak crustal field regions (figure 3, right). Detectable aurora ($> 2 R$ in clear coherent patches) are detectable in approximately 67% of observations, are more common in the dusk versus dawn quadrants, and exhibit significant variability over timescales of ~ 20 minutes, reflecting natural variability in electron dynamics and magnetic field topology in the Mars near-space environment.

Looking forward: These synoptic images, combined with simultaneous measurements of suprathermal electrons and magnetic geometry and topology from MAVEN and Mars Express promise to elucidate the complex plasma processes driving Mars’ enigmatic aurora.

References: [1] Schneider, N. M., et al. 2015, *Science*, 350, [2] Deighan, J., et al. 2018, *Nat Astron*, 2, 802, [3] Bertaux, J. L., et al. 2005, *Nature*, 435, 790, [4] Brain, D. A., et al. 2006, *Geophysical Research Letters*, 33, [5] Xu, S., et al. 2020, *Geophysical Research Letters*, 47, e87414, [6] Schneider, N. M., et al. 2018, *Geophysical Research Letters*, 45, 7391, [7]

