

# THERMAL STRUCTURE OF THE MARTIAN THERMOSPHERE: LMD-IAA GCM AND MTGCM INTERCOMPARISONS.

**F. González-Galindo**, *Instituto de Astrofísica de Andalucía, Granada, CSIC, Spain (ggalindo@iaa.es)*, **S. Bougher**, *Department of Atmospheric, Oceanic and Space Science, University of Michigan, USA*, **M.A. López-Valverde**, *Instituto de Astrofísica de Andalucía, Granada, CSIC, Spain*, **F. Forget**, *Laboratoire de Météorologie Dynamique, Institut Pierre Simon Laplace, Paris, France*, **J. Bell**, *Department of Atmospheric, Oceanic and Space Science, University of Michigan, USA*.

## Introduction

Several General Circulation Models for the study of the Martian atmosphere have been developed in the last decades, like the NASA Ames MGCM (Pollack *et al.*, 1990; Haberle *et al.*, 1993, 1999), the thermospheric MTGCM originally developed at the National Center for Atmospheric Research (Bougher *et al.*, 1990, 1999a, 2000), the GFDL Mars-GCM (Wilson and Hamilton, 1996) and the European model developed by the Laboratoire de Météorologie Dynamique (LMD, Paris) and the Atmospheric, Oceanic and Planetary Physics department (AOPP, Oxford University), LMD-AOPP GCM (Hourdin *et al.*, 1993; Read *et al.*, 1997; Forget *et al.*, 1999). With the exception of the MTGCM, all these models have been devoted to studies of the lowest layers of the Martian atmosphere. Recently, in collaboration with the Instituto de Astrofísica de Andalucía (IAA, CSIC, Granada) the LMD model has been extended to thermospheric altitudes, becoming thus the first GCM covering the whole Martian atmosphere (Angelats i Coll *et al.*, 2005; González-Galindo *et al.*, 2005).

For these models, the validation process can include comparisons with experimental data and comparisons with results from other models. Due to the scarcity of data in the Martian mesosphere and thermosphere, the comparisons between different models are specially important for thermospheric altitudes. This is specially useful for the newly developed extension to the thermosphere of the LMD-IAA GCM. The MTGCM has already been validated against several spacecraft datasets, including those from the MGS Accelerometer & Radio Science investigations (Bougher *et al.*, 1999b; 2001; 2003; 2004; 2005) and the recent Odyssey Accelerometer experiment (Bougher *et al.*, 2005). However, the MTGCM has not been tested in detail against other Mars thermospheric models. Here we present the first inter-comparison between these two Martian thermospheric GCMs.

## Models

MTGCM is a finite difference primitive equation model that solves for time-dependent neutral temperatures, neutral-ion densities, and three-component neutral winds. Prognostic equations for the major neutral species, selected

minor species and several ions are included. These fields are simulated on 33 pressure levels above 1.32  $\mu$ bar, using a log-pressure vertical coordinate. The MTGCM is presently coupled to the NASA Ames MGCM at the 1.32  $\mu$ bar level (60-80 km). This coupling allows both migrating and non-migrating upward propagating tides to cross the MTGCM lower boundary and the effects of the thermal expansion and contraction of the Mars lower atmosphere to extend to the thermosphere. Energetic processes incorporated include solar EUV, UV, and near-IR heating, a fast non-LTE CO<sub>2</sub> 15-micron cooling parameterization (López-Valverde and López-Puertas, 2001), and molecular and eddy thermal conduction. For more details on the MTGCM code and its coupling to the NASA Ames MGCM, see Bougher *et al.*, (2004; 2005).

The LMD-IAA model, evolved from a terrestrial model and including the relevant processes near the surface, extends now from the ground up to the thermosphere (0-250 km). The radiative processes considered include the effects of dust and CO<sub>2</sub> in the lower atmosphere, as well as the absorption of UV and EUV radiation by CO<sub>2</sub>, O<sub>2</sub>, O(<sup>3</sup>P), H<sub>2</sub>, H<sub>2</sub>O and H<sub>2</sub>O<sub>2</sub>. Non-LTE effects are considered using parameterizations for the solar near-IR heating, following López-Valverde *et al.*, (1998) and for the IR cooling by CO<sub>2</sub> at 15  $\mu$ m following López-Valverde and López-Puertas, (2001). A photochemical model of the neutral atmosphere, as well as the effects of molecular diffusion, thermal conduction and a simple thermal scape scheme are also included. More details can be found in Forget *et al.*, (1999) and Angelats i Coll *et al.*, (2004; 2005).

The philosophy of this study is to keep the models essentially unchanged to see the differences between the “nominal” results. However, when some sources of discrepancy have been identified, modifications have been introduced to the models in order to study the effects of different ways of treating similar processes.

## Scenarios

Some basic scenarios have been designed for this study. The basic idea was to keep the scenarios as simple as possible, but close to real scenarios. This will allow, in case of discrepancy between both models, to use some of the existing experimental data to see which of them is closer to reality. We also wanted to explore the different




Figure 1: MTGCM zonal mean temperature structure for Ls=90

sources of variability between both models. So, scenarios for different seasons had to be designed. Keeping this in mind, we decided to use initially three different scenarios:

- Scenario #1: Ls=0, no dust
- Scenario #2: Ls=90, shallow dust distribution
- Scenario #3: Ls=270, heavy dust load

For the scenarios #2 and #3 the dust is homogeneously distributed in the horizontal, with a vertical distribution given by a Conrath distribution (Conrath, 1975) with  $\tau = 0.3$  and  $\tau = 1$  for the integrated optical dust opacity and values for the Conrath parameter of 0.3 and 0.03, respectively.

The same value for the UV and EUV heating efficiency, 18%, and for the CO<sub>2</sub>-O deactivation rate,  $3 \cdot 10^{-12} \text{ cm}^3/\text{s}$  have been used in both models. When possible, similar spectroscopic data and solar flux values have been used. By keeping the inputs to both models as similar as possible, we make sure that the differences observed are not due to different input data.

### Selection of results

A large number of maps from each model has been obtained for this comparative exercise. However, we will focus on this talk on a short selection on them, using zonal mean maps for a better display of the results.

Although this intercomparison exercise is focused on the thermal structure of the thermosphere, the temperature is not the only field that has been compared. The winds (zonal, meridional and vertical), the CO<sub>2</sub> and O number densities, and the main heating terms have also been analysed. This will be useful as a diagnostic tool,




Figure 2: LMD-IAA zonal mean temperature structure for Ls=90

helping us to understand the reasons of the differences obtained in the thermal structure.

For a better visualization of the results, we decided to use zonal mean plots of those magnitudes. These plots allow us to capture both the latitude and altitude structure of the results (see Figs. 1 and 2; detailed discussions on these plots will be given in the presentation). However, in some cases, a more detailed inspection is needed. For these cases, we also use longitude-latitude maps at three different pressure levels:  $P=10^{-3}$  Pa,  $1.2 \cdot 10^{-4}$  Pa, and  $2.4 \cdot 10^{-8}$  Pa.

Additionally, a wave decomposition of the results at some fixed altitude levels has been also made. This decomposition provides a means to quantitatively understand the effects of tides propagating from the lower atmosphere upon thermospheric structures in both models.

### Summary

The results of an intercomparison campaign between the only two Martian thermospheric GCMs operative nowadays will be shown. This will be a valuable exercise for both models, showing their strong and weak points. This will be useful to identify future improvements to be incorporated. For the LMD-IAA GCM this is only the beginning of a more extensive validation process, including comparisons with data from past and present Martian missions.

### References

Angelats i Coll, M., F. Forget, M.A. López-Valverde and F. González-Galindo (2005), The first Mars thermospheric general circulation model: the Martian atmo-

sphere from the ground to 240 km, *Geophys. Res. Lett.*, 32, L04201, doi:10.1029/2004GL021368

Angelats i Coll, M., F. Forget, M.A. López-Valverde, P.L. Read and S. Lewis (2004), Upper atmosphere of Mars up to 120 km: Mars Global Surveyor accelerometer data analysis with the LMD general circulation model, *J. Geophys. Res.*, 109, doi:10.1029/2003JE002163

Bouguer, S. W., J. R. Murphy, J. M. Bell, M. A. Lopez-Valverde, and P. G. Withers (2005), Polar warming in the Mars lower thermosphere :Seasonal variations owing to changing insolation and dust distributions, *Geophys. Res. Lett.*, in press.

Bouguer, S. W., S. Engel, D. P. Hinson, and J. R. Murphy (2004) ,MGS Radio Science Electron Density Profiles : Interannual variability and implications for the neutral atmosphere, *J. Geophys. Res.*, 109, E3, E03010, doi:10.1029/2003JE002154

Bouguer, S. W., S. Engel, and P. G. Withers, (2003) The NCAR Mars Thermospheric General Circulation Model : A Review, Published Conference Abstract, International Workshop:Mars Atmosphere Modeling and Observations, Granada, Spain, January 13-15.

Bouguer, S. W., D. P. Hinson, J. M. Forbes, and S. Engel (2001) ,MGS Radio Science Electron Density Profiles and Implications for the Neutral Atmosphere, *Geophys. Res. Lett.*, 28, 3091-3094.

Bouguer, S. W., S. Engel, R. G. Roble, and B. Foster (2000), Comparative Terrestrial Planet Thermospheres : 3. Solar Cycle Variation of Global Structure and Winds at Solstices, *J. Geophys. Res.*, 105, 17669-17689.

Bouguer, S. W., S. Engel, R. G. Roble, and B. Foster (1999a), Comparative Terrestrial Planet Thermospheres : 2. Solar Cycle Variation of Global Structure and Winds at Equinox, *J. Geophys. Res.*, 104, 16591-16611.

Bouguer, S. W., G. M. Keating, R. W. Zurek, J. M. Murphy, R. M. Haberle, J. Hollingsworth, and R. T. Clancy (1999b), Mars Global Surveyor Aerobraking: Atmospheric Trends and Model Interpretation, *Adv. in Space Research*, 23, #11, 1887-1897.

Bouguer, S. W., R. G. Roble, E. C. Ridley, and R. E. Dickinson (1990), The Mars thermosphere II. General circulation with coupled dynamics and composition, *J. Geophys. Res.*, 95, 14811-14827.

Forget, F., F. Hourdin, R. Fournier, C. Hourdin and O. Talagrand (1999), Improved general circulation mod-

els of the Martian atmosphere from the surface to above 80 km, *J. Geophys. Res.*, 104, 24155-24175.

González-Galindo, F., M.A. López-Valverde, M. Angelats i Coll and F. Forget (2005), Extension of a Martian general circulation model to thermospheric altitudes: UV heating and photochemical models, *J. Geophys. Res.*, 110, E09008, doi:10.1029/2004JE002312.

Haberle, R. M., M. M. Joshi, J. R. Murphy, J. R. Barnes, J. T. Schofield, G. Wilson, M. A. López-Valverde, J. L. Hollingsworth, A. F. C. Bridger and J. Schaeffer (1999), General circulation model simulations of the Mars Pathfinder atmospheric structure investigation/meteorology data, *J. Geophys. Res.*, 104, 8957-8974.

Haberle, R. M., J. B. Pollack, J. R. Barnes, R. W. Zurek, C. B. Leovy, J. R. Murphy, H. Lee and J. Schaeffer (1993), Mars atmospheric dynamics as simulated by the NASA Ames General Circulation Model, 1, The zonal-mean circulation, *J. Geophys. Res.*, 98, 3093-3123.

Hourdin, F., P. Le Van, F. Forget and O. Talagrand (1993), Meteorological variability and the annual surface pressure cycle on Mars, *J. Atmos. Sci.*, 50, 3625-3640.

López-Valverde, M. A. and M. López-Puertas (2001), A fast computation of radiative heating rates under non-LTE in a CO<sub>2</sub> atmosphere. In *IRS 2000: Current Problems in Atmospheric Radiation*, Ed. Smith and Timofeyev, A. Deepak Publishing, Hampton, Virginia.

López-Valverde, M. A., D. P. Edwards, M. López-Puertas and C. Roldán (1998), Non-local thermodynamic equilibrium in general circulation models of the Martian atmosphere 1. Effects of the local thermodynamic equilibrium approximation on thermal cooling and solar heating, *J. Geophys. Res.*, 103, 16799-16811.

Pollack, J.B., R. M. Haberle, J. Schaeffer and H. Lee (1990), Simulations of the general circulation of the Martian atmosphere, 1, Polar processes, *J. Geophys. Res.*, 95, 1447-1474.

Read, P.L., M. Collins, F. Forget, R. Fournier, F. Hourdin, S.R. Lewis, O. Talagrand, F.W. Taylor and N.P.J. Thomas (1997), A GCM climate database for Mars: for mission planning and for scientific studies, *Adv. Space Res.*, 19 pp. 1213-1222.

Wilson, R.J. and K. Hamilton (1996), Comprehensive model simulations of thermal tides in the Martian atmosphere, *J. Atmos. Sci.*, 53, 1290-1326